

Completing Artin's braid group on infinitely many strands

Paul Fabel

Department of Mathematics & Statistics
Mississippi State University

Abstract

A generalization of the topological fundamental group is developed in order to construct a completion of Artin's braid group on infinitely many strands with respect to the following notion of convergence: $b_n \rightarrow id$ iff for each $M > 0$, eventually the first M strands of b_n are trivial.

1 Introduction

Artin's braid group B_n is the fundamental group of the collection of planar sets each of which has n elements. Fixing a basepoint $Z_n \subset R^2$, elements of B_n can be seen as collections of disjoint arcs in $R^2 \times [0, 1]$ connecting $Z_n \times \{0\}$ to $Z_n \times \{1\}$.

Artin's braid group on infinitely many strands B_∞ is the direct limit of B_n under monomorphisms $i_n : B_n \rightarrow B_{n+1}$ that attach a 'trivial strand'. It has been studied from various perspectives by Broto [2], Dehornoy [3] [4], Dynnikov [5], Moran [8], Kim and Rolfsen [10], Vershinin [11] and others.

The group B_∞ becomes a *topological* group under the requirement that a sequence of braids $b_n \rightarrow id \in B_\infty$ iff for each M there exists N such that if $n \geq N$ then the first M strands of b_n are trivial.

Unfortunately B_∞ admits no complete metric compatible with its topology. The goal of this paper is to enlarge B_∞ to a complete group of braids on infinitely many strands.

Recent work of Biss [1] shows generally that the based fundamental group of a space $\pi_1(X, p)$ inherits a canonical topology from X and becomes a topological group.

In an effort order to complete B_∞ , we develop a generalization of the topological fundamental group and subsequently exhibit a completely

metrizable topological group $\overline{B_\infty}$ such that B_∞ is embedded in $\overline{B_\infty}$ as a dense subgroup. By construction each element of $\overline{B_\infty}$ can be seen as a braid with infinitely many strands, a countable collection of pairwise disjoint arcs in $R^2 \times [0, 1]$.

The bulk of this paper is devoted to proving, for a certain space X (the space of injections of the integers into the plane), that the topological fundamental group of X is isomorphic to $\lim_{\leftarrow} G_n$, the inverse limit of Artin's pure braid groups on n strands. This serves to complete Artin's *pure* braid group on infinitely many strands, the subgroup of braids where each strand has identical endpoints. A more general construction is then employed to handle complications created by permutation of the strand endpoints.

The construction of $\overline{B_\infty}$ in this paper serves as a counterpart to that of [6] in which B_∞ is completed into a group of tame braids via the mapping class group of a disk with infinitely many punctures. However the topology of B_∞ in [6] is finer than that of the current paper. In [6] a sequence of braids b_n converges to *id* iff the first M strands are eventually trivial *and unlinked* from the remaining strands. The latter notion of convergence in B_∞ is compatible with a left order topology of Dehornoy [3], and the construction in [6] seems to resolve the issue of its completion as suggested by Dehornoy in [4].

By contrast, with the coarser topology of B_∞ adopted in the current paper, some elements of $\overline{B_\infty}$ are 'wild' in a strong sense: each representative of a wild element is not the restriction of an ambient isotopy of the plane.

2 Preliminaries

Spaces of continuous functions will have the compact open topology, product spaces will have the product topology, and quotient spaces will have the quotient topology.

Given topological spaces A and Y let $C(A, Y)$ denote the continuous functions from A into Y . Let $I(A, Y) = \{f \in C(A, Y) \mid f \text{ is one to one}\}$. Note elements of $I(A, Y)$ are not necessarily embeddings. Let $E(A, Y) = \{f \in C(A, Y) \mid f \text{ is a surjection}\}$. Let $H(A, A)$ denote the group of homeomorphisms from A onto A .

A surjective map $q : A \rightarrow Y$ is a **quotient map** provided V is open in Y if and only if $q^{-1}(V)$ is open in A . The following propositions (Theorem 11.1 p. 139 and Corollary 5.4 p287) are proved in Munkres [9].

Proposition 1 *If $q : A \rightarrow Y$ is a quotient map and $f : A \rightarrow B$ a map such that f is constant on sets of the form $q^{-1}(y)$ then there exists a*

map $F : Y \rightarrow B$ such that $f = F(q)$.

Proposition 2 *If X is locally compact and Hausdorff then $F : X \times Z \rightarrow Y$ is continuous iff $\hat{F} : Z \rightarrow C(X, Y)$ is continuous where $\hat{F}$ is defined by $(\hat{F}(z))(x) = F(x, z)$.*

If $A_1, A_2, \dots$ is a sequence of sets and $\psi_n : A_n \rightarrow A_{n-1}$ is a surjection then the **inverse limit** of A_n under ψ_n is denoted $\lim_{\leftarrow} A_n$ and consists of all sequences $(a_1, a_2, \dots)$ such that $a_n \in A_n$ and $\psi_n(a_n) = a_{n-1}$.

If the topological space G is a group with multiplication m and inversion v then G is a **topological group** provided m and v are continuous.

Let $Z^+ = \{(n, 0) \in R^2 | n \in \{1, 2, 3, \dots\}\}$.

Let $Z_n = \{(k, 0) \in Z^+ | 1 \leq k \leq n\}$.

Lemma 3 *The map $\pi : H(R^2, R^2) \rightarrow I(Z_n, R^2)$ defined via $\pi(h) = h|_{Z_n}$ determines a fibre bundle.*

Proof. Given x in the interior of a closed round planar disk D there is a circle's worth of line segments each of which has one endpoint x and the other endpoint on ∂D . Given $\{x, y\} \subset \text{int}(D)$ there is a canonical homeomorphism $h(D, x, y) : D \rightarrow D$ mapping x to y , mapping one segment linearly onto another, and fixing pointwise ∂D . Given $\alpha \in I(Z_n, R^2)$ choose h_α such that $\pi(h_\alpha) = \alpha = h_{\alpha|_{Z_n}}$. Choose disjoint closed disks D_i centered at $\alpha(i)$. For β close to α let $h_{\alpha, \beta} : R^2 \rightarrow R^2$ map $\alpha(i)$ to $\beta(i)$, leaving invariant D_i via the canonical homeomorphism, and fixing pointwise $R^2 \setminus \bigcup D_i$. Let $F = \{h \in H(R^2, R^2) | h|_{Z_n} = \text{id}|_{Z_n}\}$. For a small neighborhood U of α define $\phi : U \times F \rightarrow H(R^2, R^2)$ by $\phi(\beta, f) = (h_{\alpha, \beta})h_\alpha f$. ■

Lemma 4 *Suppose $[\alpha_0] = [\text{id}] \in \pi_1(I(Z_n, R^2), \text{id}|_{Z_n})$ and α_t is a path homotopy from α_0 to id in $I(Z_n, R^2)$. Then there exists a level preserving isotopy $h_t : R^2 \times [0, 1] \rightarrow R^2 \times [0, 1]$ such that $h_t|_{Z_n} = \alpha_t$, $h_1 = \text{id}|_{R^2 \times [0, 1]}$, and h_t is the identity on $R^2 \times \{0\}$ and $R^2 \times \{1\}$.*

Proof. Define a map $q : [0, 1] \times [0, 1] \rightarrow I(F, \text{int}(D))$ via $q(s, t) = \alpha_t(s)$. Note $\text{id}|_F = q(0, t) = q(s, 1) = q(1, t)$. The space obtained by collapsing 3 consecutive sides of $[0, 1] \times [0, 1]$ to a point is a topologically a disk. Thus by Proposition 3 q lifts to a map $Q : [0, 1] \times [0, 1] \rightarrow H(R^2, R^2)$ such that $\pi(Q) = q$ and $\text{id}|_D = Q(0, t) = Q(s, 1) = Q(1, t)$. Define $h_t : R^2 \times [0, 1] \rightarrow R^2 \times [0, 1]$ via $h_t(z, s) = Q(s, t)(z)$. ■

Lemma 5 *If $Z = \{1, 2, 3, \dots\}$ then $H(Z, Z)$ is a completely metrizable topological group.*

Proof. For $\{n, m\} \subset Z$ let $d^*(n, m) = 1$ if $n \neq m$ and $d^*(n, m) = 0$ if $n = m$. Note $C(Z, Z)$ is complete under the metric $d(f, g) = \sum_{n=1}^{\infty} \frac{d^*(f(n), g(n))}{2^n}$. Let $E^n \subset C(Z, Z)$ denote the maps whose image contains $\{1, \dots, n\}$. Suppose $f \in E^n$. Choose finite $M \subset Z$ such that $\{1, \dots, n\} \subset f(M)$ and $k = f(m_k)$ with $1 \leq k \leq n$ and $m_k \in M$. Note for g sufficiently close to f we have $g(m_k) = f(m_k)$. Hence E^n is open in $C(Z, Z)$. Thus $E(Z, Z)$ is a G_δ subspace of $C(Z, Z)$ since $E(Z, Z) = \cap E^n$. In particular $E(Z, Z)$ is completely metrizable (p.270 [9]). Let $I^n \subset C(Z, Z)$ denote the maps f such that $f|_{\{1, \dots, n\}}$ is one to one. Suppose $f \notin I^n$. Then for g sufficiently close to f we have $f|_{\{1, \dots, n\}} = g|_{\{1, \dots, n\}}$. Hence I^n is closed in $C(Z, Z)$. Thus $I(Z, Z)$ is closed in $C(Z, Z)$ since $I(Z, Z) = \cap I^n$. Note $H(Z, Z) = I(Z, Z) \cap E(Z, Z)$. Hence $H(Z, Z)$ is completely metrizable since it's a closed subspace of the completely metrizable space $E(Z, Z)$.

To prove $H(Z, Z)$ is a topological group note first that compact sets in Z are exactly the finite subsets of Z . Hence convergence in $H(Z, Z)$ is equivalent to pointwise convergence. If $f_n \rightarrow f$ and $g_n \rightarrow g$ then $\lim_{n \rightarrow \infty} f_n(g_n)(m) \rightarrow f(g)(m)$. Thus function composition is continuous in $H(Z, Z)$. To check inversion is continuous it suffices to check the case when $h_n \rightarrow id$. Fixing i choose N such that $h_n(i) = i$ whenever $n \geq N$. In particular $h_n^{-1}(i) = i$. Thus $h_n^{-1} \rightarrow id$, and hence $H(Z, Z)$ is a topological group. ■

Lemma 6 B_∞ is topologically incomplete with the topology described in the introduction.

Proof. Note that B_∞ is countable and that the complement of every one point subset of B_∞ is open and dense. Thus B_∞ is the countable union of closed nowhere dense sets. Hence by the Baire category theorem B_∞ does not admit a complete metric. ■

2.1 Generalizing the topological fundamental group

It is unknown whether there exists a space whose topological fundamental group completes B_∞ with the topology described in the introduction. To bypass this difficulty we develop a generalization of the topological fundamental group. The basic idea is to replace the basepoint of X with a subset $A \subset X$, and to consider paths of embeddings of A into X that initially fix A pointwise, that are allowed to move A off of itself in midjourney, and that place A back onto a permutation of itself at the last moment. Under suitably nice conditions the path homotopy classes determine a topological group.

Suppose throughout this section that $A \subset Y$, $\{id_A\} \subset X \subset I(A, Y)$, and H is a subgroup of $H(A, A)$.

Let $K = \{f \in C([0, 1], X) \mid f(0) = id_A, f(1) \in H\}$.

Define $m : K \times K \rightarrow K$ such that

$$m(f, g)(t) = \begin{cases} f(2t) & t \leq 1/2 \\ g(2t - 1)f(1) & 1/2 \leq t \leq 1 \end{cases}$$

Define $v : K \rightarrow K$ such that $(vf)(t) = f(1 - t)(f(1))^{-1}$ whenever $f \in K$.

Endowed with the quotient topology, let $G(X, H)$ denote the space whose elements are path components of K .

Theorem 7 *If H is a topological group then m and v induce a topological group structure on $G(X, H)$.*

Proof. Evidently m is continuous and v is continuous since inversion in H is continuous. Given $f \in K$, let $[f]$ denote the path component of f in K . Let $q : K \rightarrow G(X, H)$ be the quotient map $q(f) = [f]$. Define $M : G(X, H) \times G(X, H) \rightarrow G(X, H)$ such that $M([f], [g]) = [m(f, g)]$. Define $V : G(X, H) \rightarrow G(X, H)$ via $V([f]) = [v(f)]$. To check that M and V are well defined and continuous it suffices by Proposition 1 to note that $q(m)$ and $q(j)$ are constant respectively on sets $(q \times q)^{-1}([f], [g])$ and $q^{-1}([f])$. Define $ID : [0, 1] \rightarrow K$ via $ID(t) = id_A$. The proof that $G(X, H)$ is a group with identity $[ID]$, multiplication M , and inversion V is similar to the proof that the familiar fundamental group is indeed a group. Note the paths $m(ID, f)$, $m(f, ID)$ and f all differ by an order preserving surjective map of $[0, 1]$ and hence $M([ID], [f]) = M([f], [ID]) = [f]$. Similarly the maps $m(m(f, g), h)$ and $m(f, m(g, h))$ differ by an order preserving surjective map of $[0, 1]$ and thus M is associative. Note $m(f, v(f)) = m(v(f), f)$. The path $m(f, v(f))$ can be deformed to ID by concatenating $f_{[s, 0]}$ with $f_{[s, 0]}$. Thus V serves as inversion and hence $G(X, H)$ is a topological group. ■

The based topological fundamental group $\pi_1(Y, p)$ [1] is the quotient space of loops based at p from $[0, 1]$ into Y under the equivalence relation path homotopy. This can be seen as a special case of the generalized topological fundamental group $G(X, H)$.

Corollary 8 *If $A = \{p\} \subset Y$, $H = \{id_A\}$ and $X = I(\{p\}, Y)$ then $G(X, H) \cong \pi_1(Y, p)$, the topological fundamental group of X based at p .*

2.2 Twisted Products and Inverse Limit Spaces

Lemma 12 helps establish that the inverse limit of the groups G_n is canonically isomorphic (and homeomorphic) to an iterated semidirect product of groups K_n .

Definition 9 Suppose each of $G_1, G_2, \dots, K_1, K_2, \dots$ is a group such that for all $n \geq 1$ $\cup_{i=1}^n K_i \subset G_n$ and $K_n \triangleleft G_n$. Then $\Pi_{n=1}^\infty K_n$ forms a group under the binary operation determined coordinatewise by

$$[(x_1, x_2, \dots) * (y_1, y_2, \dots)]_n = (y_1 \dots y_{n-1})^{-1} (x_n) (y_1 \dots y_{n-1}) y_n$$

(with multiplication performed in G_n). Call this group $\Pi_{G_n} K_n$, the **twisted product** of K_n over G_n .

Lemma 10 The twisted product is a well defined group.

Proof. Note $*$ is well defined since $K_n \triangleleft G_n$ and $K_i \subset G_n$ for $1 \leq i < n$. Checking associativity,

$$\begin{aligned} [x * (y * z)]_n &= [(y_1 z_1)(z_1^{-1} y_2 z_1 z_2)((z_1 z_2)^{-1} y_3(z_1 z_2 z_3)) \dots \\ &((z_1 \dots z_{n-2})^{-1} y_{n-1} z_1 z_2 \dots z_{n-1})]^{-1} x_n [(y_1 z_1)(z_1^{-1} y_2 z_1 z_2)((z_1 z_2)^{-1} y_3(z_1 z_2 z_3)) \dots \\ &((z_1 \dots z_{n-2})^{-1} y_{n-1} z_1 z_2 \dots z_{n-1})] (z_1 \dots z_{n-1})^{-1} y_n (z_1 \dots z_n) = \\ &(z_1 \dots z_{n-1})^{-1} (y_1 \dots y_{n-1})^{-1} x_n (y_1 \dots y_n) (z_1 \dots z_n) = [(x * y) * z]_n. \end{aligned}$$

Let $ID = (id, id, \dots)$. Then $[IDx]_n = x_n = [xID]_n$. Finally given $y = (y_1, y_2, \dots)$ let $x = (x_1, x_2, \dots)$ with $x_n = (y_1 \dots y_{n-1})(y_1 \dots y_{n-1} y_n)^{-1}$. Then

$$[x * y]_n = (y_1 \dots y_{n-1})^{-1} (y_1 \dots y_{n-1}) (y_n)^{-1} (y_1 \dots y_{n-1})^{-1} (y_1 \dots y_n) = id.$$

■

Lemma 11 Suppose G is a group with subgroups $\{id\} = G_0 \subset G_1 \subset G_2 \subset G_3 \dots \subset G$ and there exist homomorphisms $\phi_n : G \rightarrow G_n$ for all $n \geq 0$ satisfying $\phi_n|_{G_n} = id|_{G_n}$. Define for all $n \geq 1$ $\psi_n : G_n \rightarrow G_{n-1}$ as $\psi_n = \phi_{n-1}|_{G_n}$. Let $K_n = \ker \psi_n$. Then $\phi : G \rightarrow \Pi_{G_n} K_i$ defined as $(\phi(g))_n = (\phi_{n-1}(g))^{-1} \phi_n(g)$ is a homomorphism.

Proof. Note $K_n = \ker \psi_n \triangleleft G_n$ and $K_i \subset G_i \subset G_n$ for $1 \leq i < n$ and the hypothesis of Definition 9 is satisfied. To check $im(\phi) \subset \Pi K_i$ we verify $(\phi_{n-1}(g))^{-1} \phi_n(g) \in K_n$. Let $p = \phi_n(g)$. Note $\phi_{n-1}(g) = \phi_{n-1}(\phi_n(g)) = \phi_{n-1}(p)$ and $\phi_{n-1}(\phi_{n-1}(x)) = \phi_{n-1}(x)$ for all $x \in G$. Thus

$$\psi_n((\phi_{n-1}(p))^{-1} p) = \phi_{n-1}(\phi_{n-1}(p^{-1})) \phi_{n-1}(p) = (\phi_{n-1}(p^{-1})) \phi_{n-1}(p) = id.$$

To check ϕ is a homomorphism use the following notation. For $g \in G$ let $g_n = (\phi(g))_n \in K_n$ and let $g^{(n)} = \phi_n(g) \in G_n$. Note

$$g^{(n)} = (g^{(0)})^{-1} (g^{(1)}) (g^{(1)})^{-1} \dots (g^{(n-1)})^{-1} g^{(n)} = g_1 g_2 \dots g_n.$$

Thus

$$\begin{aligned} (\phi(gh))_n &= (\phi_{n-1}(gh))^{-1}\phi_n(gh) = \phi_{n-1}(h^{-1})(\phi_{n-1}(g^{-1})\phi_n(g)\phi_n(h) = \\ &= (h^{(n-1)})^{-1}g_nh^{(n)} = (h_1..h_{n-1})g_n(h_1...h_{n-1})h_n = (\phi(g) * \phi(h))_n. \end{aligned}$$

■

Lemma 12 Suppose $\{id\} = G_0 \subset G_1 \subset G_2..$ is a sequence of groups and $\psi_n : G_n \rightarrow G_{n-1}$ is an epimorphism such that $\psi_n|_{G_{n-1}} = id_{G_{n-1}}$. Let $\lim_{\leftarrow} G_n = \{(g_1, g_2, ..) | g_n \in G_n \text{ and } \psi_n(g_n) = g_{n-1}\}$ with group operation $(g_1, g_2, ..) * (h_1, h_2, ..) = (g_1h_1, g_2h_2, ..)$. Let $K_n = \ker \psi_n$. Then $\psi : \Pi_{G_n} K_n \rightarrow \lim_{\leftarrow} G_n$ defined as $\psi(k_1, k_2, ..) = (k_1, k_1k_2, ..)$ is an isomorphism. If for all n G_n has the discrete topology and $\lim_{\leftarrow} G_n$ and $\Pi_{G_n} K_n$ have the product topology then ψ is a homeomorphism.

Proof. To check ψ is a homomorphism

$$\begin{aligned} \psi[(x_1, x_2, ...)(y_1, y_2, ..)] &= \phi(x_1y_1, (y_1)^{-1}x_2y_1y_2, ...) = \\ x_1y_1, x_1x_2y_1y_2, ... &= (x_1, x_1x_2, ..) * (y_1, y_1y_2, ..) = \psi(x_1, x_2, ..) * \psi(y_1, y_2, ..). \end{aligned}$$

Suppose $\psi(k_1, k_2, ..) = (id, id, ..)$. Then $k_1 = id$, $k_1k_2 = id$, ... Hence by induction $k_n = id$ for all n . Suppose $(g_1, g_2, ..) \in \lim_{\leftarrow} G_n$. Let $k_1 = g_1$. Suppose by induction that $g_{n-1} = k_1...k_{n-1}$ with $k_i \in K_i$. Let $k_n = (g_{n-1})^{-1}g_n$.

Note

$$\psi_n(k_n) = \psi_n((g_{n-1})^{-1}\psi_n(g_n)) = g_{n-1}^{-1}\psi(g_n) = \psi(g_n^{-1})\psi(g_n) = id.$$

Thus ψ is an isomorphism. Note $\lim_{\leftarrow} G_n$ and $\Pi_{G_n} K_n$ are metrizable since each of G_n and K_n is metrizable. Thus it suffices to prove the homomorphisms ψ and ψ^{-1} preserve convergence at $(id, id, ..)$. Suppose

$$(k_1^1, k_2^1, k_3^1, ...), (k_1^2, k_2^2, k_3^2, ..), ... \rightarrow (id, id, id, ..).$$

Suppose $N > 0$. Choose M so that $k_j^i = id$ for $i \geq M$ and $j \leq N$. Suppose $i \geq M$ then $[\psi(k_1^i, k_2^i, ..)]_n = id$. Thus ψ is continuous. Conversely suppose

$$(g_1^1, g_2^1, g_3^1, ..), (g_1^2, g_2^2, g_3^2, ..), ... \rightarrow (id, id, id, ...).$$

Suppose $N > 0$. Choose M so that $i \geq M$ and $g_N^i = g_{N-1}^i = id$. Then $k_n = id$. Thus ψ^{-1} is continuous. ■

3 The complete braid groups G and $\overline{B_\infty}$ on infinitely many strands

Let $X = I(Z^+, R^2)$

Let $X_n = \{f \in X \mid f(i, 0) = (i, 0) \text{ for all } i > n \text{ and for all } i \leq n \text{ } f(i, 0) \in (-\infty, n + 2/3] \times R\}$.

Recalling Theorem 7, let $G = G(X, \{id_{Z^+}\})$, the topological fundamental group of X based at id_{Z^+} .

Viewed as a collection of arcs in $R^2 \times [0, 1]$, given $g \in [g] \in G$, the strands of g connect $(n, 0, 0)$ to $(n, 0, 1)$ for $n \in \{1, 2, 3, \dots\}$.

The main goal is to prove that G is isomorphic to the inverse $\lim_{\leftarrow} G_n$ where G_n is the pure braid group on n strands, realized as the image of G under retractions that move the first n strands ‘out of the way’ and then replace the remaining strands with trivial strands.

3.1 The retraction $P_n : X \rightarrow X_n$

Lemma 13 *There exists a retraction $P_n : X \rightarrow X_n$.*

Proof. Define $R_n : X \rightarrow [0, \infty)$ as $R_n(g) = \min\{r \mid r \geq 0 \text{ and for all } i \leq n \text{ } g(i, 0) \in (-\infty, n + 2/3 + r] \times R\}$. Define $T : R \rightarrow C(R^2, R^2)$ as $T(r)(x, y) = (x - r, y)$. Define $P_n : X \rightarrow X_n$ via $P_n(g)(i, 0) = \begin{cases} (TR_n g)(g(i, 0)) & i \leq n \\ (i, 0) & i > n \end{cases}$. ■

3.2 The groups G_n and homomorphisms ϕ_n and ϕ

Suppose $n \geq 1$. Define $P_n^* : \pi_1(X, id) \rightarrow \pi_1(X_n, id)$ as $P_n^*[g] = [P_n(g)]$ where $P_n : X \rightarrow X_n$ is the retraction from Lemma 13. Let $i_n^* : \pi_1(X_n, id) \rightarrow \pi_1(X, id)$ satisfy $i_n^*[\alpha] = [\alpha]$. Recall $G = \pi_1(X, id_{Z^+})$. Let $G_n = im(i_n^*) \subset G$. Define $\phi_n : G \rightarrow G_n$ as $\phi_n = i_n^* P_n^*$. Let $G_0 = \{[id]\} \subset G$ and define $\phi_0 : G \rightarrow G_0$. Let $K_n = G_n \cap \ker \phi_{n-1}$. Define $\phi : G \rightarrow \Pi_{G_n} K_n$ as in Lemma 11.

3.3 $\phi : G \rightarrow \Pi_{G_n} K_n$ is onto

Lemma 14 *Suppose $n \geq 1$ and $[k_n] \in K_n$. Then there exists $\beta \in [k_n]$ such that $\beta(s)(i, 0) = (i, 0)$ for all $s \in [0, 1]$ and for all $i \leq n - 1$.*

Proof. Since $[k_n] \in im(i_n^*)$, we may choose $\alpha \in [k_n]$ such that the i th strand of α is straight for all $i > n$. The braid α can be deformed in X_n to a braid Γ so that the first $n - 1$ strands of Γ lie over the closed half plane $(\infty, n - \frac{1}{3}] \times R$ as follows. Recalling the map R_n from the proof of Lemma 13, fix t and translate $\alpha(t)(Z_n)$ leftward at constant speed by a total amount $R_{n-1}(\alpha(t))$. The braid $\gamma = P_{n-1}(\Gamma)$ is equivalent to the trivial braid in X_{n-1} and the first $n - 1$ strands of γ agree with the first $n - 1$

strands of Γ . Let γ_t be a path of braids in X_{n-1} taking γ to id . By Lemma 4 extend γ_t to a level preserving isotopy $h_t : R^2 \times [0, 1] \rightarrow R^2 \times [0, 1]$ such that $h_1 = id_{|R^2 \times [0, 1]}$ and h_t is the identity on the top and bottom. Now consider the path of braids $\Gamma_t = h_t h_0^{-1}(\Gamma)$. Note $\Gamma_0 = \Gamma$ and the first $n - 1$ strands of Γ_1 are straight. Let $\beta = \Gamma_1$. ■

Corollary 15 ϕ is an epimorphism.

Proof. Suppose $\{[k_1], [k_2], \dots\} \in \Pi_{G_n} K_i$.

Choose by Lemma 14 $\beta_n \in [k_n]$ such that $\beta_n(s)(i, 0) = (i, 0)$ for $i \leq n - 1$. Define $g : [1, \infty] \rightarrow X$ as

$$g(n + t) = \begin{cases} \beta_n(t) & n \geq 1 \text{ and } t \in [0, 1] \\ id & t = \infty \end{cases}.$$

Note g is well defined and continuous when $t < \infty$ since $\beta_n(0) = \beta_n(1) = id$ for $n \geq 1$. Suppose $t = \infty$, $T_n \rightarrow \infty$ and $F \subset Z^+$ is compact. Let $M = \max\{i | (i, 0) \in F\}$. For $T \geq M + 1$ and $(i, 0) \in F$, $g(T)(i, 0) = (i, 0)$. Hence g is continuous. Let $h : [0, 1] \rightarrow [1, \infty]$ be any order preserving homeomorphism. Let $f = gh$. Note $f(0) = f(1) = id$. Note $\phi_n([f]) = [\beta_1 \beta_2 \dots \beta_n]$. Thus, recalling Lemma 11, $(\phi([f]))_n = \phi_{n-1}([f])^{-1} \phi_n([f]) = [\beta_1 \dots \beta_{n-1}]^{-1} [\beta_1 \beta_2 \dots \beta_n] = [\beta_n]$. Hence ϕ is an epimorphism. ■

3.4 $\phi : G \rightarrow \Pi_G K_n$ is one to one

Lemma 16 says the following: Given an infinite pure braid α , if the infinite braid $P_{n+1}(\alpha)$ is equivalent to the trivial braid on infinitely many strands, then $\alpha|_{Z_{n+1}}$ is equivalent to the trivial braid on $n + 1$ strands.

Lemma 16 Suppose $[\alpha] \in \ker \phi_{n+1}$. Define $\beta : [0, 1] \rightarrow I(Z_{n+1}, R^2)$ as $\beta(s) = \alpha(s)|_{Z_{n+1}}$. Then $[\beta] = [id] \in \pi_1(I(Z_{n+1}, R^2), id)$.

Proof. For $(s, t) \in [0, 1] \times [0, 1]$ define $\beta_s^t : Z_{n+1} \hookrightarrow R^2$ via $\beta_s^t(s)(i, 0) = (T(tR_{n+1}(\alpha(s))))\alpha(i, 0)$. Note $\beta^0 = \beta$, $P_{n+1}(\alpha)|_{Z_{n+1}} = \beta^1$ and $\beta^1 \in [\beta] \in \pi_1(I(Z_{n+1}, R^2), id)$. Since $[P_{n+1}(\alpha)] = [id] \in \pi_1(X_n, id)$ there exists $\gamma : [0, 1] \rightarrow X_n$ such that $\gamma(0) = P_{n+1}(\alpha)$ and $\gamma(1) = id$. Now observe $\gamma(t)|_{Z_{n+1}}$ connects β^1 to id within $[\beta] \in \pi_1(I(Z_{n+1}, R^2), id)$. ■

Lemma 17 says the following. Suppose the pure braid on $n+1$ strands β^0 can be deformed to the trivial braid on $n + 1$ strands, and n strands of β^0 are straight. Then the deformation can be chosen to leave these n strands invariant.

Lemma 17 Suppose $[\beta^0] = [id] \in \pi_1(I(Z_{n+1}, R^2), id)$ and for all $s \in [0, 1]$ $\beta^0(s)_{Z_n} = id_{Z_n}$. Then there exists $\gamma : [0, 1] \rightarrow [\beta^0]$ such that $\gamma(0) = \beta^0$, $\gamma(1) = id$ and $\gamma(t)_{Z_n} = id_{Z_n}$.

Proof. Let $f : [0, 1] \times [0, 1] \rightarrow I(Z_{n+1}, R^2)$ satisfy $f(0, s) = \beta^0(s)$ and $f(1, s) = f(0, t) = f(1, t) = id$. Let $q(s, t) = f(s, t)_{Z_n}$. Note for $(s, t) \in \partial([0, 1] \times [0, 1])$ $q(s, t)_{Z_n} = id_{Z_n}$. Since $I(Z_n, R^2)$ is aspherical (Theorem 2[7]), viewed as a map of S^2 into $I(Z_n, R^2)$, q is inessential. By Proposition 3 the restriction map $\pi : H(R^2, R^2) \rightarrow I(Z_n, R^2)$ defined as $\pi(h) = h_{Z_n}$ determines a fibre bundle. Hence there exists a ‘lift’ $Q : [0, 1] \times [0, 1] \rightarrow H(R^2, R^2)$ such that $Q(s, t) = ID_{R^2}$ for $(s, t) \in \partial([0, 1] \times [0, 1])$ and $\pi(Q) = q$. Let $\gamma(t)(s) = (Q(s, t))^{-1}f(s, t)$. ■

Lemma 18 establishes the inductive step of our proof that ϕ is one to one.

Lemma 18 *Suppose $[\alpha_0] \in \ker \phi_{n+1}$ and for all $s \in [0, 1]$ $\alpha_0(s)|_{Z_n} = id_{Z_n}$. Then there exists a map $\Gamma : [0, 1] \rightarrow [\alpha_0]$ such that $\Gamma(0) = \alpha_0$, $\Gamma(t)(s)|_{Z_n} = id_{Z_n}$ and $\Gamma(1)(s)|_{Z_{n+1}} = id_{Z_{n+1}}$.*

Proof. Define $\beta_0 : [0, 1] \rightarrow I(Z_{n+1}, R^2)$ via $\beta_0(s) = \alpha_0(s)$ applying Lemmas 16 and 17 let β_t be a path homotopy from β_0 to id in $I(Z_{n+1}, R^2)$ such that the first n strands of β_t are straight. By Lemma 4 extend β_t to a level preserving isotopy $h_t : R^2 \times [0, 1] \rightarrow R^2 \times [0, 1]$ such that $h_1 = id_{R^2 \times [0, 1]}$ and h_t is the identity on the top and bottom level. Let $\Gamma_t = h_t h_0^{-1}(\alpha_0)$. ■

Lemma 19 $\phi : G \rightarrow \Pi_G K_n$ is one to one

Proof. By definition $\ker(\phi) = \cap_{i=n}^{\infty} \ker(\phi_n)$. To prove ϕ has trivial kernel we take $[g_0] \in \ker \phi$ and build a path g_t (parametrized over $[0, \infty]$) from g_0 to id within $[g_0]$ by the following procedure. By induction assume we have defined the homotopy g_t for $t \in [0, n]$ and that the first n strands of g_n are straight. Then straighten the $n+1$ strand of g_n for $t \in [n, n+1]$ via methods established in Lemmas 16, 17, and 18. Define $g_{\infty}(s) = id_{Z^+}$. We must prove the function $q : [0, \infty] \rightarrow C([0, 1], C(Z^+, R^2))$ defined via $q(t) = g_t$ is continuous when $t = \infty$. By Proposition 2 it suffices to show that the following map $Q : [0, \infty] \times [0, 1] \times Z^+ \rightarrow R^2$ defined via $Q(t, s, (n, 0)) = g_t(s)(n, 0)$ is continuous. Continuity is immediate if $t < \infty$. Suppose $t_k \rightarrow \infty$ and $s_k \rightarrow s$ and $(n_k, 0) \rightarrow (n, 0)$. Suppose $\varepsilon > 0$. Choose K such that if $k \geq K$ then $n_k = n$ and $t_k > n+1$. Suppose $k \geq K$. Then $Q(t_k, s_k, n_k) = (n, 0) = Q(\infty, s, n)$. Thus q is continuous. ■

3.5 G is the inverse limit of the pure braid groups

Theorem 20 $\phi : G \rightarrow \Pi_{G_n} K_n$ is a homeomorphism and G is topologically complete.

Proof. To prove $\phi_n : G \rightarrow G_n$ is continuous apply Proposition 1 to the retraction $P_n : X \rightarrow X_n$. Thus the isomorphism $H^* : G \rightarrow \lim_{\leftarrow} G_n$ defined via $H^*([g]) = (\phi_1([g]), \phi_2([g]), \dots)$ is continuous. Note $\phi = \psi H^*$ where $\psi : \lim_{\leftarrow} G_n \rightarrow \Pi_{G_n} K_n$ is the homeomorphism from Lemma 12. Thus ϕ is continuous. To check ϕ^{-1} is continuous it suffices, since $\Pi_{G_n} K_n$ is metrizable and ϕ^{-1} is a homomorphism, to show ϕ^{-1} preserves convergence at $([id], [id], \dots)$. Suppose $([k_1^1], [k_2^1], \dots), ([k_1^2], [k_2^2], \dots), \dots \rightarrow ([id], [id], \dots) \in \Pi_{G_n} K_n$. Choose $g^i \in \phi^{-1}([k_1^i], [k_2^i], \dots)$ such that (considering $im(g^i)$ as a subspace of $R^2 \times [0, 1]$) a maximal number of initial strands of $im(g)$ are straight line segments. It follows that $g^i \rightarrow id^*$ in $C([0, 1], X)$ where $id^*(t) = id_{Z^+}$.

Hence $[g^i] \rightarrow [id^*] \in G$ and thus ϕ^{-1} is continuous. Moreover G is completely metrizable since G is homeomorphic to the countable product of complete (discrete) spaces K_n . ■

3.6 The complete braid group $\overline{B_\infty}$

Recalling Theorem 7 let $\overline{B_\infty} = G(X, H(Z^+, Z^+))$.

In the definition of $G(X, H(Z^+, Z^+))$, when defining K , for convenience replace $[0, 1]$ with $[0, \infty]$ the one point compactification of $[0, \infty)$.

Elements of $\overline{B_\infty}$ are path components of K , and the graph of each element of K can be seen as a collection of arcs in $R^2 \times [0, \infty]$ connecting $Z^+ \times \{0\}$ to $Z^+ \times \{\infty\}$. Roughly speaking, two such graphs are equivalent if one can be deformed into the other while leaving the endpoints fixed.

We show $\overline{B_\infty}$ is topologically complete by exhibiting a homeomorphism onto the product $G \times H(Z^+, Z^+)$.

Lemma 21 *Define $\Pi : \overline{B_\infty} \rightarrow H(Z^+, Z^+)$ such that $\Pi([b])(i) = b(\infty)(i)$. Suppose there exists a map $\sigma : H(Z^+, Z^+) \rightarrow \overline{B_\infty}$ such that $\Pi\sigma = id$. Then $\overline{B_\infty}$ is homeomorphic to the product $G(X, id_{Z^+}) \times H(Z^+, Z^+)$.*

Proof. Define $T : K \rightarrow H(Z^+, Z^+)$ via $T(k) = k(\infty)$. Note all path components of $H(Z^+, Z^+)$ are trivial and thus T induces the map Π by Proposition 1. Moreover Π is a homomorphism and $\ker \Pi$ is precisely $G(X, id_{Z^+})$, the fundamental group of X based at id_{Z^+} . Now consider the map $F : \ker(\Pi) \times H(Z^+, Z^+) \rightarrow \overline{B_\infty}$ defined via $F(g, \tau) = g * \sigma(\tau)$. Consider the map $P : \overline{B_\infty} \rightarrow \ker(\Pi) \times H(Z^+, Z^+)$ defined via $P(b) = (b * (\sigma\Pi b)^{-1}, \Pi b)$.

To check that $b * (\sigma\Pi b)^{-1} \in \ker(\Pi)$ note $\Pi(b * (\sigma\Pi b)^{-1}) = \Pi(b) * \Pi((\sigma\Pi b)^{-1}) = \Pi(b) * [\Pi(\sigma\Pi b)]^{-1} = \Pi(b) * [\Pi b]^{-1} = id$. Finally note $FP = id$. Thus each of F and P is a homeomorphism. ■

Let $Z = \{1, 2, 3, \dots\}$ with the discrete topology. We will show that $H(Z, Z)$ is a closed subspace of $E(Z, Z) \subset C(Z, Z)$, that $E(Z, Z)$ is a

G_δ subspace of $C(Z, Z)$, and that $C(Z, Z)$ admits a complete metric. It follows that $H(Z, Z)$ admits a complete metric.

We now construct a ‘normal form’ for each permutation $\tau \in H(Z, Z)$ as an ‘infinite word’ $\sigma_{1m_1}\sigma_{2m_2}\dots$ where $\sigma_{nm} \in \sigma_\infty$ has compact support.

For $1 \leq n \leq m \in Z^+$ define $\sigma_{nm} : Z^+ \rightarrow Z^+$ via

$$\sigma_{nm}(i) = \begin{cases} i+1 & n \leq i \leq m-1 \\ n & i = m \\ i & \text{otherwise} \end{cases}$$

Define $b_{nm} : [0, 1] \rightarrow X$ such that b_{nm} is continuous and

$$\begin{cases} b_{nm}(1)(i, 0) = \sigma_{nm}(i) & \text{for all } i \\ b_{nm}(0)(i, 0) = (i, 0) & \text{for all } i \\ n-1 < |b_{nm}(t)(i, 0)| < m+1 & n \leq i \leq m \text{ and } n < m \\ b_{nm}(t)(i, 0) = (i, 0) & \text{otherwise} \end{cases}$$

Lemma 22 *For each $\tau \in H(Z, Z)$ there exists a sequence*

$$\sigma_{1m_1}, \sigma_{2m_2}, \dots \in H(Z, Z)$$

such that for all $k \geq 1$

$$(\tau^{-1})|_{\{1, \dots, k\}} = (\sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1})|_{\{1, \dots, k\}}^{-1}.$$

Proof. Suppose $\tau \in H(Z, Z)$. Let $m_1 = \tau^{-1}(1)$. Proceeding by induction suppose

$$(\tau^{-1})|_{\{1, \dots, k\}} = (\sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1})|_{\{1, \dots, k\}}^{-1}.$$

Define $m_{k+1} = \sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1} \tau^{-1}(k+1)$. Since $\sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1}$ maps $\tau^{-1}\{1, \dots, k\}$ onto $\{1, \dots, k\}$ it follows that $m_{k+1} \geq k+1$. By the induction hypothesis if $i \leq k$ then $(\sigma_{k+1m_{k+1}} \sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1})^{-1}(i) = \sigma_{1m_1}^{-1} \dots \sigma_{km_k}^{-1}(i) = \tau^{-1}(i)$ since $\sigma_{k+1m_{k+1}}$ fixes $\{1, \dots, k\}$ pointwise. If $i = k+1$ then

$$\begin{aligned} & (\sigma_{k+1m_{k+1}} \sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1})^{-1}(k+1) = \\ & \sigma_{1m_1}^{-1} \dots \sigma_{km_k}^{-1} \sigma_{k+1m_{k+1}}^{-1}(k+1) = \sigma_{1m_1}^{-1} \dots \sigma_{km_k}^{-1}(m_{k+1}) = \\ & \sigma_{1m_1}^{-1} \dots \sigma_{km_k}^{-1}(\sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1} \tau^{-1}(k+1)) = \tau^{-1}(k+1). \end{aligned}$$

■

Lemma 23 *There exists a continuous function $\sigma : H(Z, Z) \rightarrow \overline{B_\infty}$ such that $\sigma(\tau)(1)(i, 0) = (\tau(i), 0)$ for all $i \in H(Z, Z)$.*

Proof. Define $\hat{\sigma}: H(Z, Z) \rightarrow K \subset C([0, \infty], X)$ as follows.

Given $\tau \in H(Z, Z)$ choose $\sigma_{1m_1}, \sigma_{2m_2}, \dots$ via Lemma 22 so that for all $k \geq 1$ $(\tau^{-1})|_{\{1, \dots, k\}} = (\sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1})|_{\{1, \dots, k\}}^{-1}$. Define $\alpha_\tau: [0, \infty] \rightarrow X$ such that

$$\alpha_\tau(t)(i, 0) = \begin{cases} (i, 0) & t = 0 \\ b_{km_k}(t)\alpha_\tau(k-1)(i, 0) & t \in (k-1, k] \\ (\tau(i), 0) & t = \infty \end{cases}$$

Note α_τ is continuous at k since $b_{km_k}(0) = id$. To check continuity at $t = \infty$. Suppose $(i, 0) \in Z^+$. Let $(n, 0) = (\tau(i), 0)$. Suppose $t \in (k, k+1]$ and $n+1 \leq k$. Then $\alpha_\tau(k+t)(i, 0) = b_{km_k}(t)b_{k-1m_{k-1}} \dots b_{1m_1}(1)(i, 0) = b_{km_k}(t)(\sigma_{k-1m_{k-1}} \dots \sigma_{1m_1}(i), 0) = b_{km_k}(t)(n, 0) = (n, 0)$. Thus α_τ is continuous on $[0, \infty]$.

To check $\hat{\sigma}$ is continuous suppose $\tau_m \rightarrow \tau \in H(Z, Z)$.

Choose $\sigma_{1m_1}, \sigma_{2m_2}, \dots$ via Lemma 22 so that for all $k \geq 1$ $(\tau^{-1})|_{\{1, \dots, k\}} = (\sigma_{km_k} \dots \sigma_{2m_2} \sigma_{1m_1})|_{\{1, \dots, k\}}^{-1}$. Suppose $(i, 0) \in Z^+$. Let $n = \tau(i)$. Choose $M > n$ so that if $m \geq M$ then $\sigma_{km_k}^m = \sigma_{km_k}$ where $\sigma_{1m_1}^m, \sigma_{2m_2}^m \dots$ is chosen for τ_m as in Lemma 22. Suppose $m \geq M$ and $t \in [0, \infty]$. Note $\alpha_{\tau_m}(t)(i, 0) = \alpha_\tau(t)(i, 0)$. Thus $\hat{\sigma}$ is continuous.

Appealing to Proposition 1 let $\sigma: \Sigma_\infty \rightarrow \overline{B_\infty}$ be the induced map defined via $\sigma(\tau) = [\hat{\sigma}(\tau)]$. ■

Theorem 24 $\overline{B_\infty}$ is topologically complete.

Proof. Applying Lemmas 21, 5, 23 and Theorem 20 we obtain a homeomorphism between $\overline{B_\infty}$ and the product of topologically complete spaces $G(X, id_{Z^+}) \times H(Z^+, Z^+)$. ■

3.7 A geometric description of convergence in $\overline{B_\infty}$.

We indicate why convergence $b_n \rightarrow id \in \overline{B_\infty}$ is characterized by the statement ‘For each M there exists N such that if $n \geq N$ then the first M strands of b_n are trivial’. Recalling the proof of Lemma 21 note $b_n \rightarrow id$ iff $\Pi(b_n) \rightarrow id \in H(Z^+, Z^+)$ and $b_n * (\sigma \Pi b_n)^{-1} \rightarrow id \in G$. In $H(Z^+, Z^+)$ convergence is pointwise. Thus $\Pi(b_n) \rightarrow id$ iff for each M there exists N such that $n \geq N$ $\Pi(b_n)(i) = i$ for $i \leq M$ iff (recalling the definition of σ from Lemma 23) for each M there exists N such that the first M strands of $\sigma(\Pi b_n)$ are trivial. In G , convergence to id is determined by pointwise convergence to id of the coordinates in $\Pi_{G_n} K_n$ as shown in Theorem 20.

3.8 There exist wild braids in $\overline{B_\infty}$

Elements of the finite braid groups B_n can be understood as path homotopy classes of homeomorphisms of the plane which start at the identity, and place the finite set F_n onto itself at the last moment of the isotopy. However $\overline{B_\infty}$ cannot be understood in corresponding fashion due to the existence of ‘wild’ braids.

By definition elements of the pure braid subgroup $G \subset \overline{B_\infty}$ can be understood as equivalence classes of paths of embeddings of Z^+ into the plane which start and stop at id_{Z^+} . The set Z^+ is allowed to move off itself in midjourney. Consider for $t \in [0, 1]$ the following isotopy g_t of Z^+ . For $t \in [1 - \frac{1}{n}, 1 - \frac{1}{n+1}]$ let $(n, 0)$ perform a small clockwise orbit once around $(n + 1, 0)$ while leaving all other points of Z^+ fixed. Let S denote a simple closed curve bounding a convex disk D such that $\{(1, 0), (2, 0)\} \subset \text{int}(D)$ and $\emptyset = D \cap \{(3, 0), (4, 0), \dots\}$. For $t \in [0, 1]$ if we extend the isotopy g_t to include S we notice the diameter of S is forcibly stretched by arbitrarily large amounts. Hence the isotopy cannot be extended to S for $t \in [0, 1]$ and hence the braid determined by g_t is not determined by an ambient isotopy of the plane.

References

- [1] Biss, Daniel K., The topological fundamental group and generalized covering spaces. *Topology Appl.* 124 (2002), no. 3, 355–371.
- [2] Broto, K.; Vershinin, V. V., On the generalized homology of Artin groups. (Russian) *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 266 (2000), Teor. Predst. Din. Sist. Komb. i Algoritm. Metody. 5, 7–12, 336.
- [3] Dehornoy, Patrick, Braid groups and left distributive operations. *Trans. Amer. Math. Soc.* 345 (1994), no. 1, 115–150.
- [4] Dehornoy, Patrick, Transfinite braids and left distributive operations. *Math. Z.* 228 (1998), no. 3, 405–433.
- [5] Dynnikov, I. A., Finitely presented groups and semigroups in knot theory. (Russian) *Tr. Mat. Inst. Steklova* 231 (2000), Din. Sist., Avtom. i Beskon. Gruppy, 231–248; translation in *Proc. Steklov Inst. Math.* 2000, no. 4 (231), 220–237.
- [6] Fabel, Paul, The Mapping class group of a disk with infinitely many holes. Preprint <http://front.math.ucdavis.edu/math.GT/0303042>
- [7] Fadell, Edward; Van Buskirk, James, On the braid groups of E^2 and S^2 . *Bull. Amer. Math. Soc.* 67 1961 211–213.
- [8] Moran, Siegfried, Cable knots and infinite necklaces of knots. *Bull. Austral. Math. Soc.* 51 (1995), no. 1, 17–31.
- [9] Munkres, James R., *Topology: a first course*. Prentice-Hall, Inc.,

Englewood Cliffs, N.J., 1975.

- [10] Kim, Djun Maximilian; Rolfsen, Dale, An ordering for groups of pure braids and fibre-type hyperplane arrangements. *Canad. J. Math.* 55 (2003), no. 4, 822–838.
- [11] Vershinin, V. V., Braid groups and loop spaces. (Russian) *Uspekhi Mat. Nauk* 54 (1999), no. 2(326), 3–84; translation in *Russian Math. Section*